

How is the Diamagnetic Effect (DME) Relevant to Stellar Surface Phenomena?

Netzach Farbiash* and Raphael Steinitz*

**Ben-Gurion University, Beer-Sheva 84105, Israel*

Abstract. The structure of stellar atmospheres can be modified if they include embedded diverging magnetic fields. This is due to the presence of speed filters. Here we examine some of the relevant effects of speed filters, using numerical simulations of the motion of charged particles. We introduce the concept of "floating" particles and point out the relevance of the diamagnetic effect to the evolution of Maxwellian velocity distributions into non-Maxwellian distributions.

Keywords: Diamagnetic Effect, Charged Particle Acceleration

PACS: 95.30.Qd, 96.50.Pw

INTRODUCTION

The motion of charged particles in the presence of magnetic fields is a fundamental problem in astrophysics. Magnetic fields play significant roles in stellar atmospheres as well as in outer space. Diverging (geometrically) magnetic fields are common in the universe, so that it is essential to understand the consequences of the motion of a charged particle in such fields when also embedded in a gravitational field.

For the sake of simplicity as well as to obtain analytic solution, we mimic here the true magnetic field structures by using monopoles. For sufficiently small regions, a monopole field is a reasonable approximation to the true field (for example, field lines of a magnetic dipole, can be locally presented as field lines of a magnetic monopole (see fig 1)).




FIGURE 1. Dipole field approximated locally by a magnetic monopole.

SPEED FILTERS AND FLOATING PARTICLES

We follow the motion of a charged particle, located at a height z_0 above a magnetic monopole in the presence of a gravitational field. The height z_0 is chosen so that the vertical component of the Lorentz force is equal and opposite to the gravitational force ($\sum F_z = 0$). This choice ensures that the particle is "floating" (see fig. 2). The initial conditions:

$$|F_{Lorentz}| \sin \alpha = mg \quad (1)$$

$$|F_{Lorentz}| \cos \alpha = \frac{mv_{\perp}^2}{r} \quad (2)$$

From the geometry of the problem, we obtain that

$$\tan \alpha = \frac{r}{z_0},$$

and from eqs. 1 and 2, we get

$$\tan \alpha = \frac{v_{\perp}^2}{gr}.$$

Thus,

$$z_0 = \frac{v_{\perp}^2}{g}. \quad (3)$$




FIGURE 2. Speed filter.

We see from eq. (3) that at a height z_0 above the monopole, the "floating height" depends on $v_{\perp}^2$. If the particle has a velocity component parallel to component to the field, it will oscillate around z_0 . For $v_{\perp}^2 < gz_0$ the particle falls and the diamagnetic effect (DME) increases until the vertical component of the Lorentz force is equal to the

gravitational force. In this case, we have $z_{down} < z_0$. For $v_{\perp}^2 > gz_0$ the particle rises and the DME decreases until the vertical component of the Lorentz force is equal to the gravitational force. In that case, we have $z_{up} > z_0$. We conclude that the combination of the DME and gravitation can operate as a speed filter.

NUMERICAL SIMULATION

The magnetic monopole field is

$$\vec{B} = \frac{\beta}{R^3} \vec{R}, \quad (4)$$

where β is a constant.

The equation of motion of a particle (mass m , charge q) in the presence of gravitation is

$$\ddot{\vec{R}} = \left(\frac{q\beta}{m} \right) \frac{1}{R^3} (\dot{\vec{R}} \times \vec{R}) + \vec{g}. \quad (5)$$

We follow the position of a charged particle (proton) as function of time, in the presence of gravitation alone as well as in the presence of both gravitation and the DME. The particle's initial position is chosen to be at a given height above the magnetic monopole.




FIGURE 3. Height of proton as a function of time with both diamagnetic effect and gravity (red line), and with gravity only (blue line). Initial conditions are: $V_{0x} = 0.5\sqrt{3Kb \cdot T/m_p}$, $V_{0y} = 0.5\sqrt{3Kb \cdot T/m_p}$, $V_{0z} = 0$, $Z_0 = 3 \cdot 10^3 m$, and $T = 5 \cdot 10^3 K$

RESULTS

The above results are depicted in Figs. 3 and 4, while in Fig. 5, the acceleration is also shown. In the presence of the DME as well as gravitation, particles fall more slowly than




FIGURE 4. Height of proton as a function of time. Initial conditions, are the same for all graphs, except that of the temperature: $V_{0x} = 0.5\sqrt{3Kb \cdot T/m_p}$, $V_{0y} = 0.5\sqrt{3Kb \cdot T/m_p}$, $V_{0z} = 0$, $Z_0 = 3 \cdot 10^3 m$, and $T = 5 \cdot 10^3 K$. The first frame (top left) shows height of protons as function of time in the presence of gravity alone.




FIGURE 5. Acceleration along z axis of a proton as a function of time with diamagnetic effect and gravity. Initial conditions are: $V_{0x} = 0.5\sqrt{3Kb \cdot T/m_p}$, $V_{0y} = 0.5\sqrt{3Kb \cdot T/m_p}$, $V_{0z} = 0$, $Z_0 = 3 \cdot 10^3 m$, and $T = 5 \cdot 10^3 K$

in a gravitation field alone or can even rise (Fig. 3). We see from Fig. 5, that the effect of the DME increases in importance as $v_{\perp}^2$ (i.e. temperature) grows, as expected.

CONCLUSIONS

The simulations demonstrate that the hydrostatic structure of an atmosphere is modified by the DME. The values of pressure and temperature change, essentially because the

"effective" gravity is diminished (or inverted) selectively by the faster particles of the normal distribution. One can expect significant changes in the ionization equilibrium. The restructuring of the atmosphere can cause "apparent peculiarities" in the spectra of stars. The relevance of the DME to stellar atmospheric structure is, thus, due to the modification of effective gravity, which is particle speed dependence. The essence of the process is the speed filter.

In the future we will include collisions in order to follow the evolution of a Maxwellian distribution into a non-Maxwellian one, as required for the initial conditions in the theory discussed by Scudder (1992a, 1992b, 1994), and Anderson (1994).

We thank S. Owocki for useful discussions and helpful suggestions.

REFERENCES

1. S. W. Anderson, *ApJ*, **437**, 860-866 (1994)
2. J. D. Scudder, *ApJ*, **398**, 299-319 (1992a)
3. J. D. Scudder, *ApJ*, **398**, 319-349 (1992b)
4. J. D. Scudder, *ApJ*, **427**, 446-452 (1994)